Research

17 β -Estradiol is necessary for extinction of cocaine seeking in female rats

Robert C. Twining, Jennifer J. Tuscher, Elizabeth M. Doncheck, Karyn M. Frick, and Devin Mueller¹

Department of Psychology, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin 53211, USA

Human and preclinical models of addiction demonstrate that gonadal hormones modulate acquisition of drug seeking. Little is known, however, about the effects of these hormones on extinction of drug-seeking behavior. Here, we investigated how 17 β -estradiol (E₂) affects expression and extinction of cocaine seeking in female rats. Using a conditioned place preference (CPP) paradigm, ovariectomized rats were maintained throughout conditioning with 2 d of E₂ treatment followed by 2 d of vehicle treatment, or were injected with E₂ daily. Hormone injections were paired or explicitly unpaired with place conditioning sessions. Expression of a cocaine CPP was of equal magnitude regardless of conditioning protocol, suggesting that E₂ levels during conditioning did not affect subsequent CPP expression. During extinction, daily E₂ administration initially enhanced expression of the cocaine CPP, but resulted in significantly faster extinction compared to controls. Whereas E₂-treated rats were extinguished within 8 d, vehicle-treated rats maintained CPP expression for more than a month, indicative of perseveration. To determine whether E₂ could rescue extinction in these rats, half were given daily E₂ treatment and half were given vehicle. E₂-treated rats showed rapid extinction, whereas vehicle-treated rats continued to perseverate. These data demonstrate for the first time that E₂ is necessary for extinction of cocaine seeking in female rats, and that it promotes rapid extinction when administered daily. Clinically, these findings suggest that monitoring and maintaining optimal E₂ levels during exposure therapy would facilitate therapeutic interventions for female cocaine addicts.

Gonadal hormones render women more susceptible than men to developing compulsive patterns of psychostimulant use. Natural fluctuations in levels of the primary ovarian hormones, estrogens and progesterone, account for this increased abuse liability. Higher levels of estrogens, and lower levels of progesterone, are associated with greater sensitivity to the euphorogenic properties of cocaine in women (Evans 2007). This increased sensitivity may contribute to why women, compared with men, start using cocaine regularly at a younger age (Chen and Kandel 2002), transition from use to abuse more quickly (McCance-Katz et al. 1999), experience greater craving in response to drug-associated cues (Robbins et al. 1999) and stress (Potenza et al. 2012; Waldrop et al. 2012), and exhibit more severe drug-seeking behavior upon relapse (Gallop et al. 2007). In contrast, more women than men remain abstinent after treatment (Weiss et al. 1997). Thus, women are both more susceptible to cocaine abuse and, paradoxically, more responsive to treatment.

A substantial literature supports that estrogens mediate the enhancement of drug seeking observed in females (Febo et al. 2002; Larson et al. 2007; Segarra et al. 2010; Kerstetter and Kippin 2011), but little is known regarding the role of estrogens during treatment. Preclinically, treatment is modeled through extinction learning, which results in the formation of a new inhibitory memory that suppresses drug seeking (Quirk and Mueller 2008; Torregrossa and Taylor 2012). Whether estrogens are necessary for extinction of drug seeking, however, remains unknown. Extinction is a form of new learning, and the potent estrogen 17 β -estradiol (E₂) has been shown to enhance learning and memory across paradigms (Daniel 2006; Frick 2012). For example, E₂ enhances the consolidation of object recognition and spatial memory in ovariectomized female rats (Gresack and Frick 2006a) and,

importantly, higher levels of E₂ in intact female rats and humans are associated with enhanced consolidation and recall of fear extinction (Milad et al. 2009, 2010). Therefore, E₂ may be a key mediator of both acquisition and extinction of cocaine seeking in female rats. The effects of E₂ on extinction of cocaine seeking in females, however, have not yet been examined.

Here, we assessed the necessity of E₂ for extinction of cocaine seeking in female rats. Using a conditioned place preference (CPP) procedure, ovariectomized female rats were administered vehicle or E₂ during acquisition and extinction. We first determined the effects of E₂ on expression and extinction of cocaine seeking. We then investigated whether E₂ administration would rescue an extinction deficit in perseverating rats. Our results demonstrate that E₂ is necessary for extinction in female rats, and can facilitate extinction even after long-term perseveration of cocaine seeking.

Results

Paired E₂ during conditioning does not augment cocaine seeking

We first examined the effects of E₂ during conditioning on the magnitude of cocaine seeking on the first CPP test. To avoid buildup of E₂ levels across days, we used a water-soluble form of E₂ that is metabolized within 24 h (Pitha and Pitha 1985). E₂ injections (0.2 mg/kg, i.p.) were either paired (Paired) or explicitly unpaired (Unpaired) with conditioning trials (see Table 1, Experiment 1). After conditioning, rats were given 4 d with no E₂ treatment to eliminate carryover effects on the first day of CPP testing (extinction day 1). Both groups were injected with E₂ and tested 1 h later in the absence of cocaine for a cocaine-induced CPP. Both groups exhibited robust and virtually identical cocaine-induced CPPs (Fig. 1A). ANOVA revealed a significant effect of Chamber for both the Paired ($F_{(2,20)} = 12.64$, $P < 0.001$) and Unpaired groups ($F_{(2,22)} = 15.02$, $P < 0.001$), but

¹Corresponding author
E-mail: devinm@uwm.edu

Article is online at <http://www.learnmem.org/cgi/doi/10.1101/lm.030304.113>.

Table 1. Timeline and treatment conditions

	Conditioning						Extinction	
	(Days 1–2)	(Days 3–4)	(Days 5–6)	(Days 7–8)	...	(Days 15–16)	Trial 1	Trial 2+
Experiment 1								
Paired	E ₂ + Cond	Veh	E ₂ + Cond	Veh	...	Veh	E ₂	E ₂
Unpaired	E ₂	Veh + Cond	E ₂	Veh + Cond	...	Veh + Cond	E ₂	Veh
Experiment 2			(Days 1–8)				Trial 1	Trial 2+
E ₂			Daily E ₂ + Cond				E ₂	E ₂
Vehicle			Daily E ₂ + Cond				Veh	Veh

(E₂) 17 β -estradiol, (Veh) vehicle, (Cond) conditioning.

no Group $\times$ Chamber interaction ($F_{(2,42)} = 0.54$, $P = 0.59$). Post hoc tests confirmed that both groups spent significantly more time in the cocaine-paired chamber than either the saline-paired or center chambers (P 's < 0.05). Moreover, E₂ treatment during conditioning had no effect on motor behavior, as both Paired and Unpaired groups displayed equivalent levels of photobeam breaks during the initial CPP trial ($t_{(21)} = 1.96$, $P > 0.05$) (Fig. 1B). Taken together, these data indicate that levels of E₂ at the time of conditioning do not affect subsequent cocaine seeking.

E₂ enhances expression and extinction of cocaine seeking

To determine the extent to which E₂ affects extinction of cocaine seeking, rats received daily injections of vehicle or E₂ before each extinction trial. Repeated E₂ treatment enhanced expression of the cocaine CPP initially, but promoted rapid extinction relative to vehicle-treated controls (Fig. 2). A mixed model three-way ANOVA (Group $\times$ Chamber $\times$ Extinction Trial) revealed a significant three-way interaction ($F_{(28,588)} = 1.53$, $P = 0.045$), indicating that E₂-treated rats showed a greater CPP magnitude initially but faster extinction than vehicle-treated rats. Post hoc tests confirmed that E₂-treated rats exhibited a significantly greater CPP than vehicle-treated rats on trials 4 and 6 (P 's < 0.05) and a lack of CPP after trial 9 (P 's > 0.05). Vehicle-treated rats, however, exhibited a marked deficit in extinction across trials (Fig. 2), with post hoc tests confirming expression of a CPP through trial 16 (P 's < 0.05). Furthermore, expression of the CPP continued to persist through trial 25 (data not shown). Taken together, these data suggest that E₂ facilitates both expression and extinction of cocaine seeking in female rats and that the absence of E₂ prevents extinction.

Acute E₂ does not augment cocaine seeking

Although the magnitude of the initial CPP was not affected by paired or unpaired E₂ treatment during conditioning, an effect could have been masked because all rats received E₂ prior to the first extinction trial. To eliminate the possibility that acute E₂ affects expression of CPP, two groups of ovariectomized rats were injected daily with E₂ (0.2 mg/kg) during conditioning followed by 4 d with no treatment, and then one group received E₂ before the first CPP trial while the other group received vehicle (Table 1, Experiment 2). Both groups exhibited robust and virtually identical cocaine-induced CPPs (Fig. 3A), indicating that acute E₂ injection did not increase expression of a cocaine CPP. Repeated measures ANOVAs revealed a significant effect of Chamber for both groups (E₂, $F_{(2,20)} = 7.17$, $P = 0.005$; Vehicle, $F_{(2,22)} = 8.18$, $P = 0.003$), but no Group $\times$ Chamber interaction ($F_{(2,42)} = 0.12$, $P = 0.88$). Post hoc tests confirmed that both groups spent significantly more time in the cocaine-paired chamber than in either the saline-paired or center chambers (P 's < 0.05). Moreover, E₂ treatment prior to the initial CPP trial had no effect on motor behavior, as both E₂- and vehicle-treated groups displayed equivalent

amounts of photobeam breaks during the initial CPP trial ($t_{(21)} = 1.11$, $P > 0.05$) (Fig. 3B). Collectively, these data indicate that the timing of E₂ administration during conditioning does not affect CPP magnitude, and that expression of drug seeking does not depend on the acute effects of E₂ at test.

E₂ prevents perseverative cocaine seeking

Although daily E₂ treatment during extinction enhanced extinction of drug seeking relative to vehicle treatment, these effects may be altered by the shift from cycling E₂ on and off during conditioning to daily E₂ treatment during extinction. To test whether the noncycling groups (Experiment 2) exhibit similar extinction patterns to those of the cycling groups (Experiment 1) and to confirm the extent to which daily E₂ enhances extinction of a CPP, rats treated with E₂ on day 1 of extinction continued during extinction with daily injections of E₂, whereas rats treated with vehicle on day 1 of extinction continued with daily vehicle injections. As in Experiment 1, we found that daily E₂ treatment augmented cocaine seeking before rapidly enhancing its extinction (Fig. 4). ANOVA revealed a significant three-way interaction (Group $\times$ Chamber $\times$ Extinction Trial, $F_{(28,588)} = 1.80$, $P = 0.008$) and post hoc tests confirmed that E₂-treated rats exhibited a consistent preference for the cocaine-paired chamber on extinction trials 2–6 that was significantly higher than that of vehicle-treated rats on trials 4 and 5 (P 's < 0.05). Importantly, this CPP rapidly extinguished and was no longer evident by trial 7 (P 's > 0.05). Without E₂ treatment, however, the vehicle group exhibited a marked deficit in extinction learning across trials (Fig. 4). Post hoc tests confirmed this observed deficit, as these animals exhibited an inconsistent CPP

Figure 1. Paired and unpaired E₂ treatment during conditioning did not affect initial CPP expression. (A) E₂ administered 1 h before (Paired) or during the 2 d between (Unpaired) conditioning trials resulted in a similar preference for the cocaine-paired chamber compared with either the saline-paired or center chambers. (B) E₂ treatment had no effect on locomotor activity, as measured by photobeam breaks, when administered prior to a CPP trial regardless of E₂ treatment during conditioning. (*) $P < 0.05$ relative to the saline-paired and center chambers.

Figure 2. E₂-enhanced expression and extinction of cocaine seeking. Ovariectomized rats injected daily with E₂ (0.2 mg/kg, i.p.) during extinction initially expressed a larger cocaine-induced CPP than vehicle-treated rats, but this extinguished more rapidly than in vehicle-treated rats. Vehicle-treated rats initially expressed a weaker CPP and then exhibited a pronounced deficit in extinction learning that led to a larger CPP in later trials. (*) $P < 0.05$, Cocaine vs. Saline; (#*) $P < 0.05$, E₂ vs. Vehicle.

during early trials that grew more consistent as extinction training continued (P 's < 0.05). Indeed, the CPP was significantly larger than the extinguished preference of their E₂-treated counterparts on trials 9 and 16 (P 's < 0.05). Repeated testing revealed that the vehicle-treated rats failed to extinguish drug seeking across 36 daily trials, and the magnitude of the CPP increased across trials (Fig. 4). ANOVA revealed a significant two-way interaction (Chamber $\times$ Extinction Trial, $F_{(70,770)} = 1.43$, $P = 0.015$) and post hoc tests confirmed that vehicle-treated rats continued to express a significant cocaine-induced CPP on trials 17–36 (P 's < 0.05). Together, these data indicate that E₂ is sufficient to prevent perseverative cocaine seeking in ovariectomized female rats, and that a lack of E₂ replacement prevents extinction learning in this paradigm altogether.

E₂ rescues a persistent deficit in extinction of cocaine seeking

Perseveration of drug seeking is a hallmark of addiction (American Psychiatric Association, 2000), and our data show that without E₂ replacement, ovariectomized female rats persistently seek out the cocaine-paired chamber in spite of 25 additional extinction trials. Whether E₂ would facilitate extinction in these perseverating rats, however, remained unknown. Thus, we asked whether E₂ treatment could rescue extinction learning in these rats. Perseverating rats were divided into two groups receiving either vehicle or E₂ 1 h before further extinction training. We found that daily E₂ treatment rescued extinction learning within seven additional extinction trials (trials 37–43), even after 5 wk of perseverative drug seeking (Fig. 5). ANOVA revealed a significant Chamber $\times$ Trial interaction ($F_{(2,6)} = 9.93$, $P = 0.013$) between trial 36 (last trial without E₂) and the terminal trial 43 (last trial with E₂) for rats injected with E₂. Post hoc tests for E₂-treated rats confirmed that the cocaine CPP on day 36 was no longer significant by trial 43; moreover, the amount of time spent on the cocaine-paired side on trial 36 was significantly greater than that of trial 43 (P 's < 0.05). In contrast, ANOVA revealed no difference in the expression of CPP of vehicle-treated rats between trials 36 and 43 ($F_{(2,6)} = 0.62$, $P = 0.57$) (Fig. 5). Indeed, there was a significant Chamber effect on trial 43 ($F_{(2,6)} = 7.36$, $P < 0.024$), and post hoc tests confirmed that vehicle-treated rats continued to spend significantly more time in the cocaine-paired chamber than either the saline-paired or center chambers (P 's < 0.05). Taken together,

these results demonstrate that E₂ can rescue the capacity to extinguish drug seeking.

Discussion

We found that E₂ facilitates extinction of cocaine seeking in ovariectomized female rats independently of its well-characterized enhancement of cocaine's unconditioned hedonic effects (Becker and Hu 2008). The magnitude of the initial CPP was identical regardless of whether E₂ was explicitly paired or unpaired with cocaine (Experiment 1) or administered daily (Experiment 2) during conditioning. Moreover, acute E₂ treatment just prior to the first extinction trial had no effect on the magnitude of CPP expression. Therefore, the E₂-enhanced expression of cocaine seeking observed during extinction is not likely due to

E₂-mediated enhancements of either cocaine reward or learning during conditioning. Rather, our results indicate that E₂ treatment during extinction training leads to both greater expression and faster extinction through a mnemonic mechanism. Moreover, without E₂ replacement, ovariectomized female rats fail to suppress cocaine-seeking behavior under extinction conditions. Remarkably, this extinction learning deficit can be rescued by E₂ treatment even after 5 wk of perseverative cocaine seeking. As such, daily E₂ treatment is sufficient to enhance the cognitive processes necessary for extinction learning to proceed. These data suggest that although E₂ promotes cocaine intake in female rats self-administering cocaine (Zhao and Becker 2010), subsequent treatment with this hormone also enhances the learning required to suppress cocaine seeking during extinction.

The finding that E₂ enhances extinction of cocaine seeking in females is consistent with previous studies showing that E₂ enhances performance on a wide range of cognitive tasks, including extinction learning in other paradigms (Milad et al. 2009; Phan et al. 2012; Zhao et al. 2012). E₂ replacement or elevated E₂ levels are associated with improved object and social recognition (Gresack and Frick 2006a; Phan et al. 2012), spatial reference memory in a water maze (Gresack and Frick 2006a), and extinction of fear in female rats and humans (Milad et al. 2009, 2010).

Figure 3. Acute E₂ did not affect expression of a cocaine-induced CPP. (A) E₂- and vehicle-treated rats both exhibited a similar preference for the cocaine-paired chamber compared with either the saline-paired or center chambers. (B) E₂ injection had no effect on locomotor activity, as measured by photobeam breaks, when administered prior to a CPP trial. (*) $P < 0.05$ relative to the saline-paired and center chambers.

Figure 4. E₂ administered 1 h before daily extinction trials enhanced expression and extinction of a cocaine CPP in rats that were maintained on E₂ during conditioning. Ovariectomized rats injected daily with E₂ during extinction initially expressed a larger cocaine-induced CPP than vehicle-treated rats, but extinguished more rapidly than in vehicle-treated rats. Vehicle-treated rats initially expressed a weaker CPP and then exhibited a pronounced deficit in extinction learning that led to larger CPP in later trials. Vehicle-treated rats failed to extinguish drug-seeking behavior in 36 trials. (*) $P < 0.05$, Cocaine vs. Saline; (**) $P < 0.05$, E₂ vs. Vehicle on the same day.

Similarly, low estrogen levels predict resistance to extinction of fear-potentiated startle in women diagnosed with post-traumatic stress syndrome (Glover et al. 2012), and recent evidence shows that low levels of E₂ in both rats and humans leads to resistance to extinction of conditioned fear (Graham and Milad 2013). Extinction depends on new learning that suppresses the expression of the original memory (Bouton 2004). Therefore, E₂ could enhance extinction of cocaine seeking by bolstering acquisition, retrieval, and/or consolidation of the new extinction memory or, perhaps, interfering with reconsolidation of the former drug associations (Torregrossa and Taylor 2012). The latter is unlikely because E₂ would have also interfered with reconsolidation of extinction and our design employed repeated and prolonged exposure to the context, which favors the formation of extinction memories (Mamiya et al. 2009). Moreover, there is substantial evidence that E₂ treatment enhances memory consolidation in other learning paradigms (Packard and Teather 1997; Gresack and Frick 2006a), and emerging evidence that cognitive enhancers administered during the consolidation window can facilitate extinction of cocaine (Botreau et al. 2006; Paolone et al. 2009) or amphetamine CPP (Schroeder and Packard 2003). Although the mnemonic and neural mechanisms underlying E₂-induced enhancement of extinction remain unknown, the present experiments are the first to demonstrate that E₂ facilitates extinction of cocaine seeking in female rats.

Our findings suggest that E₂ is required for females to extinguish cocaine seeking. Ovariectomized female rats failed to extinguish for over a month, but extinction in these rats could be rescued with E₂ replacement. The observed perseverative drug seeking was unexpected, and indicative of the necessity of E₂ for extinction. The fact that E₂ rescues these extinction deficits suggests that E₂ enhances synaptic plasticity associated with extinction learning. Estrogen receptors α and β are expressed in the medial prefrontal cortex, amygdala, and hippocampus (Shughrue et al. 1997; Ostlund et al. 2003), structures that mediate extinction learning and consolidation (Quirk and Mueller 2008). E₂ stimulates intracellular signaling cascades, growth factor induction, synaptogenesis, and protein synthesis (Spencer et al. 2008; Frick 2012). Importantly, there is a synergistic overlap between the effects of E₂ and synaptic plasticity associated with extinction learning. For example, E₂ increases activation of mitogen-activated protein kinase (MAPK) cascades during memory consolidation (Fernandez et al. 2008), and MAPK signaling is required for extinction (Hugues et al. 2004). Thus, E₂ may promote extinction through stimulation of MAPK cascades. Similarly, levels of brain-

derived neurotrophic factor are increased following both E₂ administration (Luine and Frankfurt 2012) and successful extinction (Peters et al. 2010). Collectively, these findings suggest that E₂ acts to strengthen synaptic plasticity associated with extinction learning. As such, the absence of E₂ could suppress synaptic plasticity, thereby impairing extinction.

Our finding that E₂ transiently enhanced expression of cocaine seeking is congruent with the vast majority of literature describing the enhanced response to cocaine exhibited by E₂-treated female rats. Female rats exhibit enhanced responding to psychostimulants in multiple paradigms in a manner both dependent and independent of ovarian sex hormones (Hu et al. 2004; Jackson et al. 2006). In ovariectomized rats, E₂ enhances the sensitization of cocaine-induced locomotor behavior and CPP (Segarra et al. 2010), self-administration of cocaine on both fixed and progressive ratio schedules of reinforcement (Lynch and Taylor 2005; Zhao and Becker 2010), and preference for cocaine over food reinforcement in a discrete trials procedure (Kerstetter and Kippin 2011), and lowers thresholds for electrical brain stimulation of the ventral tegmental area (Galankin et al. 2010). Thus, E₂-enhanced expression of CPP could be interpreted as further evidence in females of "enhanced abuse liability to cocaine," yet this interpretation is at odds with the finding that E₂ also enhanced extinction of drug seeking. In fact, because E₂ augmented CPP expression only after four extinction trials, the data point

Figure 5. E₂ treatment rescues an extinction learning deficit in ovariectomized rats that perseverate on cocaine seeking. A cocaine-induced CPP that persisted for 5 wk was abolished by 7 d of E₂ administration with extinction training. Vehicle-treated rats, however, continued to express a cocaine CPP. (*) $P < 0.05$ relative to the saline-paired and center chambers.

toward an alternative explanation, specifically that a common neural and mnemonic mechanism mediates both enhanced expression and extinction. Because E_2 enhances memory in a variety of paradigms, E_2 may act to enhance working memory and/or retrieval of the original cocaine association, thereby augmenting the incentive salience of the cocaine-associated context and enhancing expression of the CPP. The same E_2 -mediated cognitive enhancement, then, would also enhance extinction learning and lead to more rapid extinction of cocaine seeking. Neurobiologically, both enhanced expression and extinction would likely be influenced by E_2 modulation of mesocortico-limbic dopamine (Bazzett and Becker 1994; Wyvell and Berridge 2001; Jacobs and D'Esposito 2011) or norepinephrine signaling (Mueller et al. 2008; Mueller and Cahill 2010; Otis and Mueller 2011). Thus, it is conceivable that previous reports demonstrating enhanced expression of cocaine-induced CPP assign too much importance to estrogen's undisputed enhancement of cocaine reward with little attention paid to the cognitive enhancements mediated by E_2 .

The present findings have clear implications for basic and clinical research, and for clinical treatment of women addicted to cocaine. Estrogens are key hormones that regulate the response to cocaine in both female rats and humans (Segarra et al. 2010). Changes in estrogen levels in cycling females may account for the apparent greater severity of cocaine addiction, or faster transition from use to abuse, relative to that of men (Becker et al. 2012). Nevertheless, more men than women initiate illicit drug use (55.1% vs. 44%; and for cocaine, 18.1% vs. 11.4%), which may account for the higher rate overall of drug abuse among men (3.6% vs. 2.0%; SAMHSA 2010). These findings suggest a differential path to abuse for men and women (for review, see Becker et al. 2012) or, alternatively, reflect the greater propensity for women to remain abstinent following treatment for drug abuse (Weiss et al. 1997). Our finding that E_2 treatment enhanced extinction of cocaine seeking even in ovariectomized rats that were perseverating for up to 5 wk suggests a potentially vital role for E_2 in the recovery from cocaine abuse. The importance of E_2 is underscored by our finding that female rats in the absence of E_2 failed to suppress cocaine seeking during extinction, and may explain why gonadally intact female rats in estrus (low E_2 levels) are more resistant to extinction of drug seeking than are male rats (Kerstetter et al. 2008; Anker et al. 2010). Therefore, low E_2 levels may increase the risk of relapse in some women, and this risk may be exacerbated by cocaine-induced menstrual cycle disorders (Mello and Mendelson 1997). Because E_2 levels may predict individual differences in response to treatment, our data suggest a potential benefit of routine assessment of hormonal status as part of treatment of cocaine addiction in women. Moreover, our findings suggest a need to maintain optimal levels of E_2 to enhance therapeutic outcome in cocaine-addicted women.

Materials and Methods

Subjects and surgery

Female Long-Evans rats (Harlan Laboratories, 175–200 g) were individually housed in clear plastic cages. Rats were maintained on a 14-h light/10-h dark cycle (lights on at 07.00 h), and had unlimited access to water and standard laboratory rat chow (Teklad, Harlan Laboratories). Rats were weighed and handled daily. All experimental protocols were approved by the Institutional Animal Care and Use Committee at the University of Wisconsin-Milwaukee in accordance with National Institutes of Health guidelines.

Rats were handled for a minimum of 3 d prior to surgery. They were then anesthetized with a mixture of ketamine/xylazine (90 mg/kg/10.5 mg/kg, i.p.), and underwent bilateral ovariecto-

my using a dorsal approach (Frick et al. 2004). Through a single incision along the spine, the ovary was isolated, the tip of the uterus was clamped and ligated, and the ovary was removed with a scalpel. The remaining tissue was returned to the abdomen, the procedure repeated on the other ovary, and the incision closed with sterile silk suture and wound clips. Following surgery, rats were given an antibiotic (penicillin G procaine, 75,000 units in 0.25 mL) and an analgesic (carprofen, 5.0 mg in 0.1 mL) subcutaneously, and were given 7 d to recover.

Drugs

Cocaine HCl (National Institute on Drug Abuse) was dissolved in sterile 0.9% saline (10 mg/mL), and administered at a dose of 10 mg/kg, i.p. To ensure that E_2 levels would not build up over time from repeated infusions, we used a water-soluble form of E_2 dissolved in 2-hydroxypropyl- β -cyclodextrin (HBC) that is metabolized within 24 h (Pitha and Pitha 1985). The HBC vehicle and HBC-encapsulated E_2 (Sigma Aldrich) were dissolved in sterile 0.9% saline (0.2 mg/mL) and injected i.p. 1 h before testing at a dose of 0.2 mg/kg. This dose produces levels of E_2 in the physiological range (Gresack and Frick 2006b). Post-training i.p. injection of this dose of HBC-encapsulated E_2 enhances spatial memory in the water maze in ovariectomized rats (Packard and Teather 1997) and mice (Gresack and Frick 2006a) and object recognition in ovariectomized mice (Gresack and Frick 2006a).

Conditioning and testing

Conditioning and testing were conducted as described previously (Otis and Mueller 2011). Briefly, a three-chamber apparatus was used in which two larger conditioning chambers were separated by a smaller center chamber. Baseline preferences were determined by placing the rats into the center chamber with free access to the entire apparatus for 15 min. Rats spent less time in the center chamber before conditioning than in either of the larger conditioning chambers ($F_{(2,94)} = 62.8$, $P < 0.001$; post hoc P 's < 0.001) and an equivalent amount of time was spent in the conditioning chambers ($P > 0.05$). Therefore, an unbiased procedure was used. Rats were randomly assigned to associate one large chamber with cocaine and the other with saline in a counterbalanced fashion over four conditioning trials. Rats were injected immediately before each 20-min conditioning session and confined to their respective chambers. For extinction training, rats were placed into the center chamber and allowed free access to the entire apparatus for 15 min. A CPP was determined when significantly more time was spent in the previously cocaine-paired chamber than in the saline-paired chamber.

Hormonal treatment and experimental design

Experiment 1

Rats were maintained on an experimenter-controlled, repeating 4-d treatment cycle consisting of 2 d of E_2 followed by 2 d of HBC vehicle (see Table 1). Conditioning trials were either paired (Paired, $n = 11$) or explicitly unpaired (Unpaired, $n = 12$) with E_2 over 16 d, resulting in four pairings with cocaine and four with saline. After conditioning, rats were given 4 d with no hormone injections before CPP testing. On the first CPP test trial, both groups were injected with E_2 and tested for a cocaine-induced CPP. For the remainder of the daily extinction trials, rats received either E_2 ($n = 11$) or HBC vehicle ($n = 12$).

Experiment 2

Rats were maintained on daily E_2 treatment throughout the conditioning phase (Table 1). Conditioning trials were conducted daily over 8 d, resulting in four pairings with cocaine and four with saline. After conditioning, rats were given 4 d with no hormone injections before CPP testing. Rats were injected with E_2 ($n = 11$) or HBC vehicle (Vehicle, $n = 12$) prior to each CPP test trial. These conditions were maintained for the remainder of extinction

testing through trial 36. To determine whether E₂ treatment could rescue extinction learning in perseverating rats, the rats treated with HBC vehicle during extinction were divided into two groups receiving either vehicle ($n = 4$) or E₂ ($n = 4$) 1 h before further extinction training. These rats received either daily E₂ treatment or vehicle for an additional seven extinction trials (trials 37–43).

Data analysis

Drug seeking during single or across multiple CPP trials was analyzed by comparing time spent in the previously cocaine-paired, saline-paired, and center chambers using multivariate mixed model and repeated measures ANOVA. When appropriate main and interaction effects were detected, Fisher's LSD post hoc tests were used to make pairwise comparisons. All data were analyzed using Statistica 64 (Statsoft, Inc.).

Acknowledgments

This research was supported by DA027870, University of Wisconsin-Milwaukee Research Growth Initiative to D.M., and University of Wisconsin-Milwaukee College of Letters & Science funding to K.M.F. We thank Patrick Reilly and Carolyn Rafa Todd for technical assistance.

References

- American Psychiatric Association. 2000. *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR)*, 4th ed, text revision. Arlington, VA. doi: 10.1176/appi.books.9780890423349.
- Anker JJ, Zlebnik NE, Carroll ME. 2010. Differential effects of allopregnanolone on the escalation of cocaine self-administration and sucrose intake in female rats. *Psychopharmacology (Berl)* **212**: 419–429.
- Bazzett TJ, Becker JB. 1994. Sex differences in the rapid and acute effects of estrogen on striatal D2 dopamine receptor binding. *Brain Res* **637**: 163–172.
- Becker JB, Hu M. 2008. Sex differences in drug abuse. *Front Neuroendocrinol* **29**: 36–47.
- Becker JB, Perry AN, Westenbroek C. 2012. Sex differences in the neural mechanisms mediating addiction: A new synthesis and hypothesis. *Biol Sex Differ* **3**: 14.
- Botreau F, Paolone G, Stewart J. 2006. D-Cycloserine facilitates extinction of a cocaine-induced conditioned place preference. *Behav Brain Res* **172**: 173–178.
- Bouton ME. 2004. Context and behavioral processes in extinction. *Learn Mem* **11**: 485–494.
- Chen K, Kandel D. 2002. Relationship between extent of cocaine use and dependence among adolescents and adults in the United States. *Drug Alcohol Depend* **68**: 65–85.
- Daniel JM. 2006. Effects of oestrogen on cognition: What have we learned from basic research? *J Neuroendocrinol* **18**: 787–795.
- Evans SM. 2007. The role of estradiol and progesterone in modulating the subjective effects of stimulants in humans. *Exp Clin Psychopharmacol* **15**: 418–426.
- Febo M, Jimenez-Rivera CA, Segarra AC. 2002. Estrogen and opioids interact to modulate the locomotor response to cocaine in the female rat. *Brain Res* **943**: 151–161.
- Fernandez SM, Lewis MC, Pechenino AS, Harburger LL, Orr PT, Gresack JE, Schafe GE, Frick KM. 2008. Estradiol-induced enhancement of object memory consolidation involves hippocampal extracellular signal-regulated kinase activation and membrane-bound estrogen receptors. *J Neurosci* **28**: 8660–8667.
- Frick KM. 2012. Building a better hormone therapy? How understanding the rapid effects of sex steroid hormones could lead to new therapeutics for age-related memory decline. *Behav Neurosci* **126**: 29–53.
- Frick KM, Fernandez SM, Bennett JC, Prange-Kiel J, MacLusky NJ, Leraneth C. 2004. Behavioral training interferes with the ability of gonadal hormones to increase CA1 spine synapse density in ovariectomized female rats. *Eur J Neurosci* **19**: 3026–3032.
- Galankin T, Shekunova E, Zvartau E. 2010. Estradiol lowers intracranial self-stimulation thresholds and enhances cocaine facilitation of intracranial self-stimulation in rats. *Horm Behav* **58**: 827–834.
- Galloo RJ, Crits-Christoph P, Ten Have TR, Barber JP, Frank A, Griffin ML, Thase ME. 2007. Differential transitions between cocaine use and abstinence for men and women. *J Consult Clin Psychol* **75**: 95–103.
- Glover EM, Jovanovic T, Mercer KB, Kerley K, Bradley B, Ressler KJ, Norrholm SD. 2012. Estrogen levels are associated with extinction deficits in women with posttraumatic stress disorder. *Biol Psychiatry* **72**: 19–24.
- Graham BM, Milad MR. 2013. Blockade of estrogen by hormonal contraceptives impairs fear extinction in female rats and women. *Biol Psychiatry* **73**: 371–378.
- Gresack JE, Frick KM. 2006a. Post-training estrogen enhances spatial and object memory consolidation in female mice. *Pharmacol Biochem Behav* **84**: 112–119.
- Gresack JE, Frick KM. 2006b. Effects of continuous and intermittent estrogen treatments on memory in aging female mice. *Brain Res* **1115**: 135–147.
- Hu M, Crombag HS, Robinson TE, Becker JB. 2004. Biological basis of sex differences in the propensity to self-administer cocaine. *Neuropsychopharmacology* **29**: 81–85.
- Hugues S, Deschaux O, Garcia R. 2004. Postextinction infusion of a mitogen-activated protein kinase inhibitor into the medial prefrontal cortex impairs memory of the extinction of conditioned fear. *Learn Mem* **11**: 540–543.
- Jackson LR, Robinson TE, Becker JB. 2006. Sex differences and hormonal influences on acquisition of cocaine self-administration in rats. *Neuropsychopharmacology* **31**: 129–138.
- Jacobs E, D'Esposito M. 2011. Estrogen shapes dopamine-dependent cognitive processes: Implications for women's health. *J Neurosci* **31**: 5286–5293.
- Kerstetter KA, Kippin TE. 2011. Impact of sex and gonadal hormones on cocaine and food reinforcement paradigms. *J Addict Res Ther* **54**: 2963.
- Kerstetter KA, Aguilar VR, Parrish AB, Kippin TE. 2008. Protracted time-dependent increases in cocaine-seeking behavior during cocaine withdrawal in female relative to male rats. *Psychopharmacology (Berl)* **198**: 63–75.
- Larson EB, Anker JJ, Gliddon LA, Fons KS, Carroll ME. 2007. Effects of estrogen and progesterone on the escalation of cocaine self-administration in female rats during extended access. *Exp Clin Psychopharmacol* **15**: 461–471.
- Luine V, Frankfurt M. 2012. Interactions between estradiol, BDNF and dendritic spines in promoting memory. *Neuroscience* doi: 10.1016/j.neuroscience.2012.10.019.
- Lynch WJ, Taylor JR. 2005. Decreased motivation following cocaine self-administration under extended access conditions: Effects of sex and ovarian hormones. *Neuropsychopharmacology* **30**: 927–935.
- Mamiya N, Fukushima H, Suzuki A, Matsuyama Z, Homma S, Frankland PW, Kida S. 2009. Brain region-specific gene expression activation required for reconsolidation and extinction of contextual fear memory. *J Neurosci* **29**: 402–413.
- McCance-Katz EF, Carroll KM, Rounsaville BJ. 1999. Gender differences in treatment-seeking cocaine abusers—implications for treatment and prognosis. *Am J Addict* **8**: 300–311.
- Mello NK, Mendelson JH. 1997. Cocaine's effects on neuroendocrine systems: Clinical and preclinical studies. *Pharmacol Biochem Behav* **57**: 571–599.
- Milad MR, Igwe SA, Lebron-Milad K, Novales JE. 2009. Estrous cycle phase and gonadal hormones influence conditioned fear extinction. *Neuroscience* **164**: 887–895.
- Milad MR, Zeidan MA, Contero A, Pitman RK, Klibanski A, Rauch SL, Goldstein JM. 2010. The influence of gonadal hormones on conditioned fear extinction in healthy humans. *Neuroscience* **168**: 652–658.
- Mueller D, Cahill SP. 2010. Noradrenergic modulation of extinction learning and exposure therapy. *Behav Brain Res* **208**: 1–11.
- Mueller D, Porter JT, Quirk GJ. 2008. Noradrenergic signaling in infralimbic cortex increases cell excitability and strengthens memory for fear extinction. *J Neurosci* **28**: 369–375.
- Ostlund H, Keller E, Hurd YL. 2003. Estrogen receptor gene expression in relation to neuropsychiatric disorders. *Ann N Y Acad Sci* **1007**: 54–63.
- Otis JM, Mueller D. 2011. Inhibition of β -adrenergic receptors induces a persistent deficit in retrieval of a cocaine-associated memory providing protection against reinstatement. *Neuropsychopharmacology* **36**: 1912–1920.
- Packard MG, Teather LA. 1997. Posttraining estradiol injections enhance memory in ovariectomized rats: Cholinergic blockade and synergism. *Neurobiol Learn Mem* **68**: 172–188.
- Paolone G, Botreau F, Stewart J. 2009. The facilitative effects of D-cycloserine on extinction of a cocaine-induced conditioned place preference can be long lasting and resistant to reinstatement. *Psychopharmacology (Berl)* **202**: 403–409.
- Peters J, Dieppa-Perea LM, Melendez LM, Quirk GJ. 2010. Induction of fear extinction with hippocampal-infralimbic BDNF. *Science* **328**: 1288–1290.

- Phan A, Gabor CS, Favaro KJ, Kaschack S, Armstrong JN, Macluskay NJ, Choleris E. 2012. Low doses of 17 β -estradiol rapidly improve learning and increase hippocampal dendritic spines. *Neuropsychopharmacology* **37**: 2299–2309.
- Pitha J, Pitha J. 1985. Amorphous water-soluble derivatives of cyclodextrins: Nontoxic dissolution enhancing excipients. *J Pharm Sci* **74**: 987–990.
- Potenza MN, Hong KI, Lacadie CM, Fulbright RK, Tuit KL, Sinha R. 2012. Neural correlates of stress-induced and cue-induced drug craving: Influences of sex and cocaine dependence. *Am J Psychiatry* **169**: 406–414.
- Quirk GJ, Mueller D. 2008. Neural mechanisms of extinction learning and retrieval. *Neuropsychopharmacology* **33**: 56–72.
- Robbins SJ, Ehrman RN, Childress AR, O'Brien CP. 1999. Comparing levels of cocaine cue reactivity in male and female outpatients. *Drug Alcohol Depend* **53**: 223–230.
- SAMHSA. 2010. National Survey on Drug Use and Health. Substance Abuse and Mental Health Services Administration: Substance Abuse and Mental Health Data Archive <http://www.icpsr.umich.edu/icpsrweb/SAMHDA/studies/32722>, doi: 10.3886/ICPSR32722.v3. ICPSR/University of Michigan Institute for Social Research, Ann Arbor, MI.
- Schroeder JP, Packard MG. 2003. Systemic or intra-amygdala injections of glucose facilitate memory consolidation for extinction of drug-induced conditioned reward. *Eur J Neurosci* **17**: 1482–1488.
- Segarra AC, Agosto-Rivera JL, Febo M, Lugo-Escobar N, Menendez-Delmestre R, Puig-Ramos A, Torres-Diaz YM. 2010. Estradiol: A key biological substrate mediating the response to cocaine in female rats. *Horm Behav* **58**: 33–43.
- Shughrue PJ, Lane MV, Merchenthaler I. 1997. Comparative distribution of estrogen receptor- α and - β mRNA in the rat central nervous system. *J Comp Neurol* **388**: 507–525.
- Spencer JL, Waters EM, Romeo RD, Wood GE, Milner TA, McEwen BS. 2008. Uncovering the mechanisms of estrogen effects on hippocampal function. *Front Neuroendocrinol* **29**: 219–237.
- Torregrassa MM, Taylor JR. 2012. Learning to forget: Manipulating extinction and reconsolidation processes to treat addiction. *Psychopharmacology (Berl)* doi: 10.1007/s00213-012-2750-9.
- Waldrop AE, Price KL, Desantis SM, Simpson AN, Back SE, McRae AL, Spratt EG, Kreek MJ, Brady KT. 2012. Community-dwelling cocaine-dependent men and women respond differently to social stressors versus cocaine cues. *Psychoneuroendocrinology* **35**: 798–806.
- Weiss RD, Martinez-Raga J, Griffin ML, Greenfield SF, Hufford C. 1997. Gender differences in cocaine dependent patients: A 6 month follow-up study. *Drug Alcohol Depend* **44**: 35–40.
- Wyvell CL, Berridge KC. 2001. Incentive sensitization by previous amphetamine exposure: Increased cue-triggered “wanting” for sucrose reward. *J Neurosci* **21**: 7831–7840.
- Zhao W, Becker JB. 2010. Sensitization enhances acquisition of cocaine self-administration in female rats: Estradiol further enhances cocaine intake after acquisition. *Horm Behav* **58**: 8–12.
- Zhao Z, Fan L, Frick KM. 2012. Epigenetic alterations regulate estradiol-induced enhancement of memory consolidation. *Proc Natl Acad Sci* **107**: 5605–5610.

Received January 9, 2013; accepted in revised form March 19, 2013.

17 β -Estradiol is necessary for extinction of cocaine seeking in female rats

Robert C. Twining, Jennifer J. Tuscher, Elizabeth M. Doncheck, et al.

Learn. Mem. 2013, **20**:

Access the most recent version at doi:[10.1101/lm.030304.113](https://doi.org/10.1101/lm.030304.113)

References

This article cites 57 articles, 8 of which can be accessed free at:
<http://learnmem.cshlp.org/content/20/6/300.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).