Learning & Memory

while it did not disrupt reaction to a new object, which is a hippocampal-independent function (Wan et al. 1999; Albasser et al. 2010). In this set of experiments, lack of effects of CaEDTA questions the real role of zinc and suggests that chelation of vesicular zinc by DEDTC would prevent co-release of both zinc and glutamate. Effects of DEDTC could thus only result from side effects of glutamate release blocking. However, Daumas et al. (2004), using contextual fear conditioning (CFC) with a single training session, with two trials spaced out by 2 min, brought conflicting results. With this rather massed training procedure, DEDTC impaired both acquisition and consolidation, which could appear as a confirmation of results obtained in MWM with massed practice. But, unexpectedly, this study showed that CaEDTA also impaired acquisition and consolidation, while a control chelator, zinc tetraacetic ethylenediamine acid (ZnEDTA), had no effect. This study showed for the first time in vivo that MF vesicular zinc released in the synaptic cleft is clearly involved in acquisition and consolidation of CFC. Zinc could thus play a direct role in this type of learning, contrary to spatial or to object location memory.

Therefore, we postulated that effects of glutamate on NMDARs located on postsynaptic elements of MF (MF-CA3), of temporo-ammonic pathway (TA-CA3), and of associative recurrent collaterals (ARCs) are sufficient to allow consolidation of spatial learning but not for the optimal consolidation of contextual learning. Indeed, in this kind of learning, direct action of zinc would be necessary. We thus expected that in spatial learning, chelation of zinc once released will be ineffective, while blocking NMDARs in the CA3 area will be enough to impair memory consolidation. On the other hand, in CFC, blocking NMDARs associated with chelation of zinc once released would be necessary to impair consolidation. To assess this hypothesis, microinjections of the NMDAR antagonist D-2-amino-5-phosphonopentanoate (AP5) alone or in association with the extracellular zinc chelator CaEDTA were done in the CA3 area after learning MWM and CFC with massed practice.

Inbred C57Bl/6J male mice, 9–12 wk old, were implanted with guide cannulae in the dorsal CA3 area as previously described (Daumas et al. 2004), then randomly allotted to an experimental group and subsequently received the same drug through the whole experimental procedure. After a week of recovery, mice underwent spatial learning with massed practice in the MWM then CFC. This testing order was chosen so that the less stressful learning task was performed first with the aim to minimize order effects. The MWM training procedure started with a familiarization session as used by Lassalle et al. (2008) except that the visible platform was moved to a different location for each familiarization trial. Massed learning with the submerged platform was realized 24 h later. Mice were given four sessions of three consecutive trials with a 15–20-min ISI. Total duration of learning procedure was limited to 90 min. Twenty-four hours after learning, mice were submitted to a 60-sec spatial test. Data collection was performed using Ethovision 2.3 (Noldus). During learning sessions, latencies to reach the platform were recorded. During the test, numbers of annuli crossings were analyzed. Statistics were performed with Systat 10.2; α levels were set at $P < 0.05$. Learning performances were analyzed using one factor repeated measures ANOVA. For spatial bias during recall, a repeated measures ANOVA was used to compare annuli crossings. Then, for each treatment, repeated measures ANOVA analyzed variation of annuli crossings. When significant changes in crossings had been detected, Fisher's LSD post-hoc test was used to determine the most frequently crossed annulus.

CFC, carried out 4 d after MWM, consisted of a single learning session with two trials as described by Daumas et al. (2004). Memory of CFC was measured 24 h later in the same experimental conditions, followed 2 h later by assessment of elementary sound–

shock association in a modified context. Freezing was used to evaluate fear memory (Blanchard and Blanchard 1969). Freezing occurrences were recorded every 5 sec according to the method of instantaneous sampling at fixed time intervals (Martin and Bateson 1986). Freezing data (p) expressed as frequency of immobility episodes were transformed as $Q = \arcsin(\sqrt{p/100})$ then processed by one factor (treatment) ANOVA. Post-hoc comparisons were made using Fisher's LSD tests.

Immediately after every learning session of MWM and CFC, mice received similar 0.25- μ L bilateral infusions of ZnEDTA (200 mM), AP5 (3.04 mM), or a mixture of AP5 and CaEDTA or AP5 and ZnEDTA in the CA3 area, as described by Daumas et al. (2004), according to their experimental group.

ZnEDTA is a chelator of divalent ions saturated with zinc. CaEDTA, a chelator of divalent ions saturated with calcium, cannot cross membranes which makes it an exclusively extracellular chelator unable to change intracellular zinc homeostasis. CaEDTA was chosen to trap specifically free zinc and zinc released by MF. Zinc affinities of CaEDTA and ZnEDTA are very high ($K_d = 10^{-16.4}$ M and $K_d = 0^{-15}$ M, respectively), whereas affinities for calcium or magnesium are very low. Consequently, CaEDTA chelates zinc without chelating calcium or magnesium while ZnEDTA chelates none, and can consequently serve as control.

AP5, a competitive antagonist of NMDARs, binds specifically on NMDARs, so preventing any connection of glutamate. The time window of CaEDTA and AP5 is estimated to be ~ 1 h (www.drugs.com/ppa/edetate-calcium-disodium-calcium-edta.html).

At the end of behavioral tests, mice were euthanized with pentobarbital (150 mg/kg, i.p., cEVA), and their brains processed for histology (Fig. 1; see also Daumas et al. 2004). Subjects with misplaced cannulae and/or injections were excluded from analyses, leading to groups of: ZnEDTA, $n = 15$; AP5, $n = 11$; ZnEDTA + AP5, $n = 14$; and CaEDTA + AP5, $n = 10$.

As can be appreciated from Figure 2A, when mice are exposed to a massed learning procedure, they gradually improve their performance. Mean escape latencies decreased significantly over trials in all groups ($F_{(3,138)} = 12.077$, $P < 0.001$), with no significant variation between groups ($F_{(3,46)} = 0.345$, $P = 0.793$). Therefore, before receiving treatment, all mice efficiently used distal visual cues and showed comparable learning profiles.

After the last training trial, animals were injected with drug depending on their experimental group, and spatial memory was assessed 24 h later. During the memory test, if mice are

Figure 1. (A) Timeline of the experimental design from surgery to sacrifice of the animals for histology. Note that each behavioral procedure consists of a day training followed by memory test on the consecutive day. (WM) Water maze, (CFC) contextual fear conditioning. (B) Photograph of a frontal section of a mouse dorsal hippocampus stained with thionin. Injection sites can be seen at the level of the CA3 area on both hippocampi (white arrows). Black arrows indicate the position of guide cannulae.

Figure 2. MWM task. (A) Learning performance: mean latency to platform for the different groups over the four massed learning sessions in the MWM. (B) Test of spatial bias: effects of four treatments (ZnEDTA, AP5, ZnEDTA + AP5, CaEDTA + AP5) on the average number of annuli crossings during the recall test at 24 h. (PF) Platform location learned. Results are presented as mean $\pm$ SEM. (*) $P < 0.05$, (***) $P < 0.001$.

remembering the previous learning correctly, they should spend more time around the platform location and therefore present a higher number of targeted annulus crossings in comparison with the three other possible positions of it. As shown in Figure 2B only control animals focused their research on the previously learned location ($P < 0.001$). None of the treated groups appropriately retrieved the correct platform location. A significant variation of search profiles between groups appeared ($F_{(3,138)} = 12.307$, $P < 0.001$), with a significant treatment $\times$ annulus crossings interaction ($F_{(9,138)} = 4661$, $P < 0.001$). Repeated measures ANOVAs performed within each group revealed a heterogeneity in patterns of annuli crossing for groups ZnEDTA ($F_{(3,42)} = 25.846$, $P < 0.001$) and AP5 ($F_{(3,30)} = 3.805$, $P = 0.020$), while groups CaEDTA + AP5 and ZnEDTA + AP5 showed no such heterogeneity ($F_{(3,39)} = 1.491$, $P = 0.232$ and $F_{(3,27)} = 0.656$, $P = 0.586$). Blocking glutamate binding on NMDARs by AP5 clearly disrupted retention of goal learning. Curiously, the AP5-alone treated group presented a peculiar profile. AP5 mice that showed no better recall of the location of the platform than the other treatment groups (AP5 + CaEDTA and AP5 + ZnEDTA) tended to always go back to the release point (opposite to PF location). They showed preferential crossings of the Opposite annulus compared to Adjacent left ($P = 0.036$) and Adjacent right annuli ($P = 0.029$). No differences appeared between search profiles of CaEDTA + AP5 and ZnEDTA + AP5 groups. Combination of extracellular zinc chelator CaEDTA with AP5 provided no additional effect compared to the combination of ZnEDTA with AP5, therefore confirming results from Lassalle et al. (2000) with distributed learning, as well as from Florian and Roulet (2004) with massed learning, about the lack of effect of CaEDTA alone.

Altogether, it can be concluded that the NMDARs antagonist (AP5) caused impaired memory consolidation in MWM, whereas the extracellular zinc chelator (CaEDTA) had no effect.

After a 4-d resting period, mice were submitted to contextual fear conditioning (CFC) before being treated with the drug of their experimental group. During conditioning, all groups show no freezing behavior before the first shock, and a progressive increase of it after the shocks ($F_{(10,440)} = 28.109$, $P < 0.001$) (data not shown).

Twenty-four hours after conditioning and treatment, contextual fear memory was first assessed (Fig. 3A). Although context re-exposure in the ZnEDTA control group resulted in slightly lower freezing levels than usually observed in our studies, possibly resulting from previous exposure to MWM learning, rates of contextual freezing showed, nevertheless, a significant variation due to treatments ($F_{(3,45)} = 9.309$, $P < 0.001$, $R^2 = 0.384$). Indeed, while the rate of freezing was 54.6% in the ZnEDTA control group, it was 27.9% in AP5, 27.4% in ZnEDTA + AP5, and 17.2% in the CaEDTA + AP5 group. Fisher's post-hoc tests confirmed significant differences of freezing rates between the ZnEDTA control

group and AP5 ($P = 0.003$), ZnEDTA + AP5 ($P = 0.002$), and CaEDTA + AP5 ($P < 0.001$) groups, respectively. Although no significant difference appeared between groups AP5 and ZnEDTA + AP5 ($P = 0.580$), confirming the lack of specific effect of ZnEDTA alone, Fisher's tests showed a significantly greater decrease of freezing rates for the CaEDTA + AP5 compared to the AP5 groups ($P = 0.032$).

Hippocampal-independent fear memory was then tested (Fig. 3B). As shown in Figure 3B, all groups express very high rates of freezing, $\sim 80\%$, during sound memory tests, clearly demonstrating that CA3-hippocampal injections did not disturb the elemental sound-shock association ($F_{(3,44)} = 0.488$, $P = 0.692$).

Therefore AP5 impaired memory consolidation of CFC while a mixture of both AP5 and CaEDTA showed a cumulative effect.

This study analyzed cumulative roles of CA3 zinc and glutamate in the consolidation stage of two forms of hippocampal-dependent memories acquired under massed learning procedures. Microinjections of AP5 in CA3 aimed at blocking NMDARs and thus postsynaptic action of glutamate at the CA3 synapses. Coinjections of CaEDTA allowed additional chelation of MF zinc without impacting glutamate release. ZnEDTA appeared to be a good control.

In the MWM, AP5 impaired memory consolidation whereas CaEDTA caused no additional effect. It can therefore be concluded there is a lack of effect of MF zinc chelation on the consolidation of spatial learning, while action of glutamate at the three synapses TA-CA3, MF-CA3, and ARCs is mandatory. On the other hand, in CFC a mixture of both AP5 and CaEDTA resulted in cumulative impairments of memory consolidation. As expected, they had no effect on memory of cue conditioning. These results confirm that consolidation of spatial and contextual memories relies on at least partially different mechanisms. However, these results are at variance with those of Fellini et al. (2009) that found no role for NMDARs in water-maze memory consolidation, using a similar

Figure 3. Effects of the four treatments (ZnEDTA, AP5, ZnEDTA + AP5, and CaEDTA + AP5) administered immediately after the conditioning and measured 24 h later, during the test to the context (A) and 26 h later during the test to the sound in the modified context (B). Results are presented as mean $\pm$ SEM. (*) $P < 0.05$, (**) $P < 0.01$, (***) $P < 0.001$.

massed procedure. This discrepancy can be explained since, in our study, during familiarization the platform was moved to a different location for each trial, whereas in the Fellini et al. (2009) study the platform remained at the same place during the whole procedure. It has been shown that using the same location for familiarization and learning results in a pre-exposure effect that contributes to the formation of a spatial representation that remains valid during the learning phase. For example, Rudy et al. (2004) have developed the concept of context pre-exposure facilitation effect (CPFE), demonstrating that when rats were pre-exposed to the conditioning context in CFC, inactivation of amygdala following conditioning was of no consequence for consolidating the memory of context because the representation was already established in the hippocampus. Similarly, Roesler et al. (1998) showed that pre-exposure of rats to the inhibitory avoidance box 24 h before training prevented the amnesic effect of AP5 hippocampal post-training infusion on NMDARs. Therefore, pre-exposure of mice to the spatial learning configuration during familiarization is likely to explain the disengagement of NMDARs during consolidation of spatial memory observed by Fellini et al. (2009) in the MWM.

Direct comparison of MWM and CFC under massed learning procedures reconciles seemingly contradictory results from our group about MF-CA3 pathway involvement in the consolidation of spatial learning in MWM, critical with massed practice (Florian and Roulet 2004) while optional with distributed practice (Lassalle et al. 2000). The importance of training procedure has been raised by Silva's group showing that mice with a partial knockout of the cAMP responsive-element binding (CREB KO) expressed almost normal long-term memories (contextual and spatial) following distributed learning practice, whereas they were severely disrupted with massed learning (Kogan et al. 1996). Therefore, by analogy, we assume that differences in memory consolidation efficiency could rely on training procedures. When MFs are selectively blocked, consolidation remains nevertheless possible in MWM with learning sessions distributed over several days. But, above all, our results strengthen previous conclusions about a specific role for zinc released into the MF-CA3 synaptic cleft on CFC consolidation (Daumas et al. 2004). Direct comparison between CFC and MWM with massed procedure shows, beyond doubt, that the action of glutamate at the three CA3 synapses is necessary and sufficient for the consolidation of spatial learning; MF-CA3 released zinc playing no direct role in this case. On the other hand, both vesicular glutamate and zinc are mandatory and exert cumulative effects on CFC memory consolidation, suggesting involvement of two complementary mechanisms in contextual learning with a strong emotional component.

Acknowledgments

This work was supported by CNRS and Université de Toulouse. We thank the ABC facility and ANEXPLO for housing mice.

References

- Albasser MM, Poirier GL, Aggleton JP. 2010. Qualitatively different modes of perirhinal-hippocampal engagement when rats explore novel vs. familiar objects as revealed by c-Fos imaging. *Eur J Neurosci* **31**: 134–147.
- Amaral DG, Witter MP. 1995. Hippocampal formation. In *The rat nervous system* (ed. Paxinos G), pp. 443–493. Academic Press, San Diego, CA.
- Assaf SY, Chung SH. 1984. Release of endogenous Zn^{2+} from brain tissue during activity. *Nature* **308**: 734–736.
- Besser L, Chorin E, Sekler I, Silverman WF, Atkin S, Russell JT, Hershfinkel M. 2009. Synaptically released zinc triggers metabotropic signaling via a zinc-sensing receptor in the hippocampus. *J Neurosci* **29**: 2890–2901.
- Blanchard RJ, Blanchard DC. 1969. Crouching as an index of fear. *J Comp Physiol Psychol* **67**: 370–375.
- Bordi F, Ugolini A. 1999. Group I metabotropic glutamate receptors: Implications for brain diseases. *Prog Neurobiol* **59**: 55–79.
- Daumas S, Halley H, Lassalle JM. 2004. Disruption of hippocampal CA3 network: Effects on episodic-like memory processing in C57BL/6J mice. *Eur J Neurosci* **20**: 597–600.
- Fellini L, Florian C, Courtney J, Roulet P. 2009. Pharmacological intervention of hippocampal CA3 NMDA receptors impairs acquisition and long-term memory retrieval of spatial pattern completion task. *Learn Mem* **16**: 387–394.
- Florian C, Roulet P. 2004. Hippocampal CA3-region is crucial for acquisition and memory consolidation in Morris water maze task in mice. *Behav Brain Res* **154**: 365–374.
- Frederickson CJ. 1989. Neurobiology of zinc and zinc-containing neurons. *Int Rev Neurobiol* **31**: 145–238.
- Kesner RP. 2007. Behavioral functions of the CA3 subregion of the hippocampus. *Learn Mem* **14**: 771–781.
- Kogan JH, Frankland PW, Blendy JA, Coblenz J, Marowitz Z, Schutz G, Silva AJ. 1996. Spaced training induces normal long-term memory in CREB mutant mice. *Curr Biol* **7**: 1–11.
- Kwon HB, Castillo PE. 2008. Long-term potentiation selectively expressed by NMDA receptors at hippocampal mossy fiber synapses. *Neuron* **57**: 108–120.
- Lassalle JM, Bataille T, Halley H. 2000. Reversible inactivation of the hippocampal mossy fiber synapses in mice impairs spatial learning, but neither consolidation nor memory retrieval, in the Morris navigation task. *Neurobiol Learn Mem* **73**: 243–257.
- Lassalle JM, Halley H, Daumas S, Verret L, Francés B. 2008. Effects of the genetic background on cognitive performances of Tg2576 mice. *Behav Brain Res* **191**: 104–110.
- Li Y, Hough CJ, Frederickson CJ, Sarvey JM. 2001a. Induction of mossy fiber → CA3 long-term potentiation requires translocation of synaptically released Zn^{2+} . *J Neurosci* **21**: 8015–8025.
- Li Y, Hough CJ, Suh SW, Sarvey JM, Frederickson CJ. 2001b. Rapid translocation of Zn^{2+} from presynaptic terminals into postsynaptic hippocampal neurons after physiological stimulation. *J Neurophysiol* **86**: 2597–2604.
- Lu YM, Taverna FA, Tu R, Ackerley CA, Wang YT, Roder J. 2000. Endogenous Zn^{2+} is required for the induction of long-term potentiation at rat hippocampal mossy fiber-CA3 synapses. *Synapse* **38**: 187–197.
- Lynch MA. 2004. Long-term potentiation and memory. *Physiol Rev* **84**: 87–136.
- Martin P, Bateson P. 1986. *Measuring behaviour: An introductory guide*. Cambridge University Press, Cambridge, UK.
- Morris RG, Moser EI, Riedel G, Martin SJ, Sandin J, Day M, O'Carroll C. 2003. Elements of a neurobiological theory of the hippocampus: The role of activity-dependent synaptic plasticity in memory. *Philos Trans R Soc Lond B Biol Sci* **358**: 773–786.
- Quinta-Ferreira ME, Matias CM, Arif M, Dionisio JC. 2004. Measurement of presynaptic zinc changes in hippocampal mossy fibers. *Brain Res* **1026**: 1–10.
- Roesler R, Vianna M, Sant'Anna M, Kuyen CR, Kruel AVS, Quevedo J, Cardoso Ferreira MB. 1998. Intrahippocampal infusion of the NMDA receptor antagonist AP5 impairs retention of an inhibitory avoidance task: Protection from impairment by pretraining or preexposure to the task apparatus. *Neurobiol Learn Mem* **69**: 87–91.
- Rolls ET. 2007. An attractor network in the hippocampus: Theory and neurophysiology. *Learn Mem* **14**: 714–731.
- Rolls ET. 2010. A computational theory of episodic memory formation in the hippocampus. *Behav Brain Res* **215**: 180–196.
- Rudy JW, Huff NC, Matus-Amat P. 2004. Understanding contextual fear conditioning: Insights from a two-process model. *Neurosci Biobehav Rev* **28**: 675–685.
- Sensi SL, Yin HZ, Carriedo SG, Rao SS, Weiss JH. 1999. Preferential Zn^{2+} influx through Ca^{2+} -permeable AMPA/kainate channels triggers prolonged mitochondrial superoxide production. *Proc Natl Acad Sci* **96**: 2414–2419.
- Stupien G, Florian C, Roulet P. 2003. Involvement of the hippocampal CA3-region in acquisition and in memory consolidation of spatial but not in object information in mice. *Neurobiol Learn Mem* **80**: 32–41.
- Takeda A, Fuke S, Minami A, Oku N. 2007. Role of zinc influx via AMPA/kainate receptor activation in metabotropic glutamate receptor-mediated calcium release. *J Neurosci Res* **85**: 1310–1317.
- Wan H, Aggleton JP, Brown MW. 1999. Different contributions of the hippocampus and perirhinal cortex to recognition memory. *J Neurosci* **19**: 1142–1148.
- Weiss JH, Sensi SL. 2000. Ca^{2+} - Zn^{2+} permeable AMPA or kainate receptors: Possible key factors in selective neurodegeneration. *Trends Neurosci* **23**: 365–371.

Received October 15, 2012; accepted in revised form April 26, 2013.

Access the most recent version at doi:[10.1101/lm.029017.112](https://doi.org/10.1101/lm.029017.112)

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).